

SUSY-induced Flavor and CP-violation in B -physics *

Gudrun Hiller
Munich (LMU-ASC)

SUSY 2005, IPPP Durham, July 23, 2005

* *“Implications of current and future b -data for (weak scale) SUSY ”*

Thanks to HFAG, Zoltan Ligeti, Guy Raz, Salim Safir for input

Non-CP observables/rare decays → plenary talk by M. Misiak (Monday)

Lepton Flavor violation → plenary talk by A. Brignole (after this talk)

SM=Standard Model, NP=New Physics, EWKSB=electroweak symmetry breaking, FCNC=Flavor changing neutral currents

SM tests with indirect processes

indirect loop processes:


$\Lambda \gtrsim m_W$ scale of New Physics

$$\mathcal{L}_{eff} = \sum_i c_i^{(n)} \frac{O_i^{(n)}}{\Lambda^n} \quad c_i^{(n)} \leftrightarrow f\left(\underbrace{m_j, g_l, \dots}_{\text{colliders}}; \underbrace{\varphi_{CKM}, \varphi_m, \delta_n}_{\text{flavorphysics}}\right)$$

no competition from large SM tree contributions

complement collider searches

FCNC: sensitivity to SM and NP phases φ and flavor-breaking couplings δ



MFV = no more flavor/CP violation than in SM, i.e. in Yukawas (CKM)

$U(3)^5$ symmetry only broken by “Y” spurion fields; RGE invariant

superpotential leads to CKM-type flavor-breaking (with unbroken R-parity)

$$-\mathcal{L}_{soft} \supset \tilde{Q}^\dagger m_Q^2 \tilde{Q} + \tilde{U}^\dagger m_U^2 \tilde{U} + \tilde{D}^\dagger m_D^2 \tilde{D} + A_U \tilde{Q} H_U \tilde{U} + A_D \tilde{Q} H_D \tilde{D}$$

squark masses (*A-terms*) : 3×3 hermitean (**complex**) matrices

super-CKM basis: quarks are mass eigenstates, but squarks not UNLESS

$$m_{Q,D,U}^2 \propto 1 \text{ and } A_{U,D} \propto Y_{U,D} \quad (1)$$

beyond (1) \rightarrow large # of new sources of flavor/CP violation


e.g. tree level FCNC $\tilde{g} \tilde{q}_i \tilde{q}_j$, mass insertions $\delta_{i,j}^{U,D} \equiv \Delta m_{i,j}^{2U,D} / m_{diag}^{2U,D}$

(1) realized in GMSB, AMSB, CMSSM (RGE induced effects tiny)

beyond MFV terms in $b \rightarrow s$ motivated from GUTs from large $\nu_\mu - \nu_\tau$ oscillations \rightarrow sizeable effects in $B_s - \bar{B}_s$ -mixing [hep-ph/0212180](https://arxiv.org/abs/hep-ph/0212180), [0212397](https://arxiv.org/abs/hep-ph/0212397)

Sensitivity to New Physics

models of EWKSB with NP @ TeV



reach in indirect signals rare b, c, K, τ -decays, mixing, EDMs, $g-2$
depends on beyond the SM flavor/CP violation (minimal=CKM ?), large
parameters e.g. $\tan\beta$ and theor. and exp. uncertainties Fig from hep-ph/0207121


- Introduction: SUSY effects in b -physics ✓
- Searching for BSM phases in $b \rightarrow s$: “ $\sin 2\beta$ ” measurements*
- Recent and future SM tests from $b \rightarrow s\gamma$, $b \rightarrow sl^+\ell^-$ processes
- Flavor violation in models beyond the minimal model
 - R-parity violating effects in double radiative decays
 - light A^0 in NMSSM at large $\tan\beta$
- Outlook

*in this talk β denotes the phase of $B\bar{B}$ -mixing, i.e. β_{eff}

Time-dependent CP asymmetries in $b \rightarrow s\bar{q}q$ decays

$$\text{SM+MFV: } -\eta_{CP} \sin 2\beta \underbrace{\left((\bar{s}s) K_S \right)}_{FCNC} = \sin 2\beta \underbrace{\left((\bar{c}c) K_S \right)}_{\text{tree}} + \underbrace{\left| \frac{V_{ub}V_{us}^*}{V_{tb}V_{ts}^*} \right|}_{\mathcal{O}(\lambda^2)} \cdot \#$$

SM background $\mathcal{O}(\lambda^2) \sim 0.04$, $\#$ non-universal, hadronic physics



since Moriond: $\sim 1\sigma$ shifts in $\bar{c}c$, Φ , larger ones in K_S^3 , f_0 ; η' off by 2.3σ
better agreement between Belle and BaBar

$$\Delta S = -\eta_f S_f - \sin 2\beta$$

hep-ph/0505075, 0503151 hep-ph/9708305, 0310020, 0303171, 0403287

f	$\Delta S_{SM}^{QCDF@NLO}$	$\Delta S_{SM}^{QCDF@LO}$	$ \Delta S_{SM}^{SU(3)+} $	S_f LP'05	C_f LP'05
ΦK^0	0.01 ... 0.03	0.02	$\lesssim 0.3$	0.47 ± 0.19	-0.09 ± 0.14
$\eta' K^0$	0 ... 0.02	-0.01 ... 0.02	$\lesssim 0.15$	0.48 ± 0.09	-0.08 ± 0.07
$\pi^0 K_S$	0.03 ... 0.12	0.03 ... 0.10	$\lesssim 0.2$	0.31 ± 0.26	-0.02 ± 0.13
ωK_S	0.05 ... 0.22	0.05 ... 0.25	-	0.63 ± 0.30	-0.44 ± 0.23


LO captures central value and uncertainty of full NLO calc. of ΔS_f

for all above modes: QCD factorization predicts $\Delta S_f > 0$, experimental shifts < 0 ; no significant $C_f \neq 0$

ultimately more precision needed; all exp. errors $\lesssim 0.1$ only by time of super- b -factory [hep-ph/0503261](#)

Pattern of NP in $b \rightarrow s$ Penguin modes

generic NP scenarios with one additional phase, use QCDF @ LO [hep-ph/0503151](https://arxiv.org/abs/hep-ph/0503151)





input: $S_{\pi^0 K_S} = 0.34$ ● and maximal NP amplitude ok with other data

predict S_f for: ▼ = Φ , ▲ = η' , ■ = η , × = ω , + = ρ

goal: identify type of NP from characteristic SM departure

Fit generic NP scenarios to current data

$A = A_{SM} + A_{NP}; A_{NP} \propto \epsilon_i e^{i\vartheta_i}$ hep-ph/0503151 χ^2 -fit to $b \rightarrow s$ penguin data



black, dark grey, light grey regions: probability $> 0.32, 0.046, 0.0027$


IFF $A_{NP} \gg A_{SM}$ 4-fold solution $\sin 2(\beta + \vartheta) \simeq 0.4$ (with $\cos 2\beta > 0$)

NP amplitudes can be larger if no phase $\vartheta = 0, \pi$

all 3 scenarios have solution, which is more favored than SM ($\epsilon_i = 0$)

Fit to SUSY-model beyond MFV

assume gluino loops are dominant: δ_{23}^D , LL, LR, RL, RR, complex
[hep-ph/0407073](https://arxiv.org/abs/hep-ph/0407073) impose bounds from $b \rightarrow s\gamma$, $b \rightarrow s\ell\ell$ (pre LP05 data)



Some (further) SM tests with b -physics

	experiment	SM	comments
$a_{CP}(b \rightarrow s\gamma)$	$0.4 \pm 3.6 \%$	$0.42 \pm 0.17 \%$ hep-ph/0312260	CPX in $bs\gamma, g$
$a_{CP}(b \rightarrow d/s\gamma)$	-0.110 ± 0.116 BaBar'05	10^{-9} hep-ph/0312260	test MFV
$S_{K_S\pi^0\gamma}$	0.00 ± 0.28 Belle/BaBar'05	$-2m_s/m_b$	V+A FCNCs
$\mathcal{B}(B \rightarrow X_s g)$	$< 9\%$ CLEO'97	$5.0 \pm 1.0 \cdot 10^{-3}$	NP in bsg
$\mathcal{B}(B \rightarrow X_s \bar{\mu}\mu)$	$4.3 \pm 1.2 \cdot 10^{-6}$	$4.3 \pm 0.7 \cdot 10^{-6}$	q^2 -spectra
$a_{CP}(B \rightarrow X_s \bar{\ell}\ell)$	-0.22 ± 0.26	$-0.2 \pm 0.2 \%$ hep-ph/9812267	CPX
$A_{FB}^{CP}(B \rightarrow K^* \bar{\ell}\ell)$	–	$\lesssim 10^{-3}$ hep-ph/0006136	CPX in bsZ
$R_K \mu\mu$ vs. ee	1.06 ± 0.48 BaBar'05	$1 + \mathcal{O}(m_\mu^2/m_b^2)$ hep-ph/0310219	non-SM Higgs
$\mathcal{B}(B \rightarrow K\nu\bar{\nu})$	$< 3.6 \cdot 10^{-5}$ Belle'05	$3.8_{-0.6}^{+1.2} \cdot 10^{-6}$	$O(10)$ from SM
$\mathcal{B}(B_s \rightarrow \mu^+\mu^-)$	$< 5.8 \cdot 10^{-7}$	$3.2 \pm 1.5 \cdot 10^{-9}$	$O(100)$ from SM
$\mathcal{B}(B_s \rightarrow \tau^+\tau^-)$	$< \mathcal{O}(5\%)$	$7.2 \pm 1.1 \cdot 10^{-7}$	$O(10^5)$ from SM
Δm_s	$> 15/\text{ps}$	$(15 - 22)/\text{ps}$	hadron colliders




Impact of $b \rightarrow sl^+l^-$ beyond MFV, perspectives

$\mathcal{B}(b \rightarrow sl^+l^-)$: best bound on Z -penguins $\bar{s}Zb$ -coupling $\sim C_{10} \lesssim 2 \times \text{SM}$
 order 1 effects in C_{10} from δ_{23}^U possible (LR and LL) [hep-ph/9906286,0006136](https://arxiv.org/abs/hep-ph/9906286)

great NP sensitivity in q^2 -spectra in $b \rightarrow sl^+l^-$; asy \neq forward - \neq
 backward l^+ in dilepton CMS w.r.t. \bar{B} **needs tagging $A_{FB}^{SM} + \bar{A}_{FB}^{SM} \simeq 0$**


$A_{FB}(\hat{s}) \sim \text{Re} \left[C_{10}^* (C_7^{eff} + \beta(\hat{s})C_9^{eff}) \right]$ also $B \rightarrow K^* l^+ l^- \rightarrow$ Belle'04

$O_{10} \sim \bar{s}_L \gamma_\mu b_L \leftrightarrow \bar{s} Z b$



shape sensitive to $\text{sign} C_7$ (no zero for $C_7 > 0$); $A_{FB} \propto C_{10}$; flat possible
zero allows precision test of SM $\hat{s}_{SM}^{NNLL} = 0.162 \pm 0.002(8)$ [hep-ph/0208088,0209006](https://arxiv.org/abs/hep-ph/0208088)


NP in $bs\gamma + \gamma$ tightly constrained by $\mathcal{B}(B \rightarrow X_s\gamma)$, but 1PI has room



impact of 4-Fermi's is higher order in $b \rightarrow s\gamma$, lepton-loop down by α_{em}

$b \rightarrow s\tau^+\tau^-$ essentially unconstrained $\mathcal{B}(B_s \rightarrow \tau^+\tau^-) < 5\%$ [hep-ph/0411344](#)

R-parity violation: biggest contribution from $\tilde{\nu}_\mu$ through $\lambda'_{232}\lambda_{233}$ [hep-ph/0404152](#)



RPV: $Br/Br_{SM}(B_s \rightarrow \gamma\gamma) \lesssim 16$ $Br/Br_{SM}(B \rightarrow X_s\gamma\gamma) \lesssim 5$


model-independently: $Br/Br_{SM}(B \rightarrow K\gamma\gamma) \lesssim O(10)$ [hep-ph/0411344](#)

$\mathcal{B}^{L3}(B_s \rightarrow \gamma\gamma) < 1.48 \cdot 10^{-4} \simeq 10^2 \cdot \mathcal{B}_{SM}(B_s \rightarrow \gamma\gamma)$

Light CP-odd A^0 : implications for hadron colliders

direct searches: $h^0 \rightarrow A^0 A^0$ open if $2m_{A^0} < m_{h^0}$

can be VERY important for $\mathcal{O}(1)$ $h^0 A^0 A^0$ -coupling c [hep-ph/0005308](https://arxiv.org/abs/hep-ph/0005308)




decay modes: $A^0 \rightarrow b\bar{b}, \tau\tau, 3\pi$ or higher hadronic, $\mu\mu, ee, \gamma\gamma$

if A^0 very light and weakly coupled, it becomes **missing energy**


bounds from Υ -decays, beam dump, astro physics $m_{A^0} \gtrsim \mathcal{O}(100\text{MeV})$ ok

NMSSM constraints from indirect signals

$W = QY_u H_u U + QY_d H_d D + LY_e H_d E + \lambda H_d H_u N - \frac{1}{3}kN^3$ N :singlet
 at large $\tan\beta$: naturally light A_1^0 , rad. stable $b \rightarrow sA_1^0$ transitions




bounds from $B \rightarrow KA_1^0$, $K \rightarrow \pi A_1^0$, $\Upsilon(1s) \rightarrow \gamma A_1^0$ decays



A_1^0 masses as low as $\mathcal{O}(10\text{MeV})$ viable [hep-ph/0404220](https://arxiv.org/abs/hep-ph/0404220)


Light CP-odd A^0 : further tests from b -physics

- improve bounds from radiative Υ -decays or $B \rightarrow K +$ missing energy
- for $m_{\psi'} < m_{A_1^0} \lesssim m_B$: search for A_1^0 in $b \rightarrow s\tau^+\tau^-$ processes
sensitivity e.g. $\mathcal{B}(B \rightarrow X_s\tau^+\tau^-) \sim 10^{-3}$



- $B_s - \bar{B}_s$ mixing and $B_s \rightarrow \mu^+\mu^-$ CDF: $\mathcal{B}(B_s \rightarrow \mu^+\mu^-) < 5.8 \cdot 10^{-7}$
in MSSM correlated [Fig adopted from hep-ph/0207241](#)

in NMSSM: SM-like Δm_s (A_1^0 contribution constrained by Δm_d), but not correlated with $\mathcal{B}(B_s \rightarrow \mu^+\mu^-)$ [hep-ph/0404220](#)



- indirect (loop) processes are unique probes of CP/flavor sector
- flavor experiments distinguish MFV vs. non-MFV; very different model-building; sensitivity to very high scales $\Lambda_{\text{SUSY}} \gg \Lambda_{\text{EWK}}$
- goal: measure ALL (flavor diagonal \rightarrow direct searches and flavor breaking \rightarrow FCNC) couplings/parameters precisely
- so far agreement with SM/CKM, “ $\sin 2\beta$ ” data moved closer to MFV some hints ($\eta' K^0$); issue exciting and unsettled \rightarrow higher precision
- many FCNC couplings only weakly or just un-constrained .. $\Delta m_s, \tau^\pm, \dots$
- scenarios beyond minimal MSSM+MFV: interplay between flavor physics and direct searches largely unexplored
- study correlations & pattern rather than individual observables

back up slides

Double radiative $b \rightarrow q\gamma\gamma$ decays, status


modes	\mathcal{B}_{SM}	90 % C.L. bounds
$B_d \rightarrow \gamma\gamma$	$3.1^{+6.4+1.0+1.0+1.3}_{-1.6-0.9-0.7-1.0} \cdot 10^{-8}$	$5.4 \cdot 10^{-6}$ Belle'05
$B_s \rightarrow \gamma\gamma$	$1.2^{+2.5+0.3+0.3+0.01}_{-0.6-0.3-0.2-0.02} \cdot 10^{-6}$	$1.48 \cdot 10^{-4}$ L3
$B \rightarrow X_s \gamma\gamma$	$(3.7 - 5.1) \cdot 10^{-7}$	—
$B \rightarrow K \gamma\gamma$	$(0.5 - 5.6) \cdot 10^{-7}$ A,B,D	—
$B \rightarrow K^* \gamma\gamma$	few $\cdot 10^{-7}$ C,D	—

A: Singer,Zhang'97 **B:** Choudhury etal'03 **C:** Choudhury etal'04 **D:** Hiller,Safir'04 (OPE for short-distance 1PR)

Note $\sum_{H=K,K^*} \mathcal{B}(B \rightarrow H\gamma\gamma) < \mathcal{B}(B \rightarrow X_s \gamma\gamma)$; semi-radiative decays suffer charmonium pollution η_c, χ ; Br 's depend on cuts

$B \rightarrow \gamma\gamma$ errors: $\lambda_B, f_B, \mu, \gamma$ Bosch,Buchalla '02

diagrams in SM



$$\mathcal{H}_{eff} = -4 \frac{G_F}{\sqrt{2}} V_{tb} V_{ts}^* \sum C_i(\mu) O_i(\mu)$$

dipole operators $O_7 \propto \bar{s}_L \sigma_{\mu\nu} b_R F^{\mu\nu}$ $O_8 \propto \bar{s}_L \sigma_{\mu\nu} b_R G^{\mu\nu}$

4-Fermi operators $O_9 \propto (\bar{s}_L \gamma_\mu b_L)(\bar{l} \gamma^\mu l)$ $O_{10} \propto (\bar{s}_L \gamma_\mu b_L)(\bar{l} \gamma^\mu \gamma_5 l)$

new physics in Wilson coefficients $C_i = C_i^{SM} + C_i^{NP}$ and/or new operators

model-independent analysis: Br 's, $A_{CP}, A_{FB} = f(C_i) \rightarrow \text{fit !}$